

Effects of R-parity Violating Couplings on CP Asymmetries in Neutral B Decays

Dafne Guetta

*Dipartimento di Fisica
Università degli Studi di Bologna
Via Irnerio 46, I-40126 Bologna, Italy*

Abstract

A detailed analysis of the effects of supersymmetric models without R-parity on various CP asymmetries in neutral B decays is given. We concentrate on models with Abelian horizontal symmetries that allow us to estimate the order of magnitude of the new effects. We focus on channels where the standard model gives clean predictions: $B_d \rightarrow \psi K_S$ and $B_d \rightarrow \phi K_S$. In the presence of new physics the two asymmetries can have a value different from $\sin 2\beta$. Moreover, they can be different from each other.

PACS number(s): 11.30.Er, 12.60.Jv, 13.20.He

E-mail address: guetta@bo.infn.it

I. INTRODUCTION

In the next few years several experiments on B -physics will take place (BaBar, BELLE, HERA B, CLEO, RUN II at FNAL) [1]. They will provide a test of many predictions of the standard model (SM). Large part of the experiments will concentrate on CP violation measurements. In this paper we study the effects of the Minimal Supersymmetric Standard Model (MSSM) without R-parity (with conserved baryon number) on specific CP asymmetries. These asymmetries can be significantly altered from their SM values if there are important new contributions to $B_d - \bar{B}_d$ mixing and/or to the decay amplitudes.

In the SM both the b -quark decays and the $B_d - \bar{B}_d$ mixing are determined by combinations of the CKM matrix elements [2]. The asymmetries measure the relative phases between these combinations. Since the phases are related to angles of the Unitarity Triangle (UT) through the CKM unitarity, a measurement of the asymmetries determines these angles.

Conversely, the SM predictions for the CP asymmetries in neutral B decays into certain CP eigenstates are determined by the values of the three angles of the UT [2]. So, their measurements will test these SM predictions and consequently provide a probe for physics beyond the SM [3].

Our work is an extension of previous studies done on how R-parity violating models can affect CP violation measurements [4–6]. We give estimates of what the models can do. In the spirit of [7] we estimate the order of magnitude of these new effects by embedding SUSY without R-parity in models with Abelian horizontal symmetries.

The paper is organized as follows. In sections II and III we give the general formalism needed to study the CP asymmetries. In section IV we briefly review SUSY models without R-parity. In section V we study the effects of R-parity violating couplings on the decay amplitudes. We consider two CP asymmetries that have equal values in the SM, but can be different in R-parity violating models. The contribution to the mixing is the same for all B_d decays, so it cancels in the difference of the two phases that can be extracted from the asymmetries. We estimate the maximal possible effects and analyze whether they are large enough to be clearly signaled out. In section VI we consider the contribution of the R-parity violating couplings to the mixing and how this affects the CP asymmetries. In section VII we estimate the effects of the R-parity violating couplings in the framework of models where the magnitude of the fermion masses and CKM mixing angles is explained by assuming some horizontal $U(1)$ symmetry. Finally section VIII contains our conclusions.

II. GENERAL FORMALISM FOR THE DEFINITION OF CP ASYMMETRY

In this section we recall the relevant formulae for the decay of neutral B mesons into CP eigenstates [2]. The time dependent CP asymmetry is defined as:

$$a_{f_{CP}}(t) \equiv \frac{\Gamma[B_{phys}^0(t) \rightarrow f_{CP}] - \Gamma[\bar{B}_{phys}^0(t) \rightarrow f_{CP}]}{\Gamma[B_{phys}^0(t) \rightarrow f_{CP}] + \Gamma[\bar{B}_{phys}^0(t) \rightarrow f_{CP}]}, \quad (2.1)$$

where B_{phys}^0 and $\bar{B}_{phys}^0$ are states tagged as pure flavour eigenstate B_d and $\bar{B}_d$ at the production.

Eq.(2.1) can be written as:

$$a_{f_{CP}}(t) \equiv a_{f_{CP}}^{\cos} \cos(\Delta M t) + a_{f_{CP}}^{\sin} \sin(\Delta M t), \quad (2.2)$$

where

$$\begin{aligned} a_{f_{CP}}^{\cos} &= \frac{(1 - |\lambda|^2)}{1 + |\lambda|^2}, \\ a_{f_{CP}}^{\sin} &= -\frac{2\text{Im}\lambda}{1 + |\lambda|^2}. \end{aligned} \quad (2.3)$$

ΔM is the mass difference between the two physical states, and

$$\lambda = \sqrt{\frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}}} \frac{\langle f_{CP} | \mathcal{H} | B_d \rangle}{\langle f_{CP} | \mathcal{H} | \bar{B}_d \rangle} = \frac{q}{p} \frac{\bar{A}}{A}, \quad (2.4)$$

where M_{12} and Γ_{12} are the non diagonal elements of the mass matrix $\mathbf{M}$ and of the matrix $\mathbf{\Gamma}$ which describes the exponential decay of the system. Since $\Gamma_{12}/M_{12} \simeq O(1\%) \ll 1$, we have

$$\left(\frac{q}{p}\right)^2 = \frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}} \simeq \frac{M_{12}^*}{M_{12}} = \exp(-4i\phi_M), \quad (2.5)$$

where ϕ_M is the $B - \bar{B}$ mixing phase.

The quantity $\text{Im}\lambda$ that can be extracted from $a_{f_{CP}}(t)$ is directly related to CKM matrix elements in the SM. For mixing in the B_d system, we have $M_{12} \propto (V_{tb}V_{td}^*)^2$ and consequently,

$$\left(\frac{q}{p}\right)_{B_d} = \frac{V_{tb}^*V_{td}}{V_{tb}V_{td}^*}. \quad (2.6)$$

As we mentioned in the introduction CP asymmetries in decays to CP eigenstates, $B_d \rightarrow f_{CP}$, provide a way to measure the three angles of the UT, defined by [2]

$$\alpha \equiv \arg\left(-\frac{V_{td}V_{tb}^*}{V_{ub}^*V_{ud}}\right) \quad \beta \equiv \arg\left(-\frac{V_{cd}V_{cb}^*}{V_{tb}^*V_{td}}\right) \quad \gamma \equiv \arg\left(-\frac{V_{ud}V_{ub}^*}{V_{cb}^*V_{cd}}\right). \quad (2.7)$$

For example, in the SM, we can measure $\sin 2\beta$ in the decay $B_d \rightarrow \psi K_S$ where we have $\bar{A}/A = [(V_{cb}V_{cs}^*)/(V_{cb}^*V_{cs})][(V_{cs}V_{cd}^*)/(V_{cs}^*V_{cd})]$, thus

$$\begin{aligned} \lambda(B \rightarrow \psi K_S) &= \left(\frac{V_{td}V_{tb}^*}{V_{td}^*V_{tb}}\right) \left(\frac{V_{cb}V_{cs}^*}{V_{cb}^*V_{cs}}\right) \left(\frac{V_{cs}V_{cd}^*}{V_{cs}^*V_{cd}}\right) \\ &\rightarrow \text{Im}\lambda = -\sin(2\beta). \end{aligned} \quad (2.8)$$

III. CP ASYMMETRIES IN NEUTRAL B DECAYS IN THE PRESENCE OF NEW PHYSICS

We study CP asymmetries in neutral B decays into final CP eigenstates. In general, a decay amplitude has contributions from several diagrams A_i with different weak phases ϕ_i

and different strong phases δ_i [2]. If, for example, there are two dominant contributions, then the amplitudes A for $B_d \rightarrow f_{CP}$ and $\bar{A}$ for $\bar{B}_d \rightarrow f_{CP}$ are described by

$$\begin{aligned} A &= A_1 e^{i\phi_1} e^{i\delta_1} + A_2 e^{i\phi_2} e^{i\delta_2} \\ \bar{A} &= A_1 e^{-i\phi_1} e^{i\delta_1} + A_2 e^{-i\phi_2} e^{i\delta_2}. \end{aligned} \quad (3.1)$$

If $A_2/A_1 = 0$ or $\phi_1 = \phi_2$, then the CP asymmetry $a_{f_{CP}}$ cleanly measures the CP violating quantity $\sin 2(\phi_M + \phi_1)$, where ϕ_M is the weak phase of the $B_d - \bar{B}_d$ mixing amplitude. We focus on cases where within the SM this is indeed the situation.

In the SM, the CP asymmetries in the decay modes $b \rightarrow c\bar{c}s$ ($B_d \rightarrow \psi K_S$), $b \rightarrow c\bar{c}d$ ($B_d \rightarrow D^+ D^-$), $b \rightarrow c\bar{u}d$ ($B_d \rightarrow D^0 \rho$) and $b \rightarrow s\bar{s}s$ ($B_d \rightarrow \phi K_S$) all measure the angle $\phi_M + \phi_1 = \beta$ in B_d decays even if they depend on different CKM factors.

From the theoretical point of view the cleanest mode is $b \rightarrow c\bar{c}s$, for which the penguin contribution, represented by A_2 , has the same phase as the tree level one, A_1 , hence $\phi_1 = \phi_2$. In this case the theoretical uncertainty for this channel is practically $\delta\phi_{SM} = 0$. This channel is the best mode also from the experimental point of view because it has a relatively large branching ratio of $O(10^{-4})$ and can be identified by clean signals: $\psi \rightarrow l^+ l^-$, $K_S \rightarrow \pi^+ \pi^-$ [1,8]. For these reasons this mode should be the reference mode to which all the other measurements are compared. Other clean modes are $b \rightarrow c\bar{u}d$ and $b \rightarrow s\bar{s}s$ which have $\delta\phi_{SM} < 0.1$ [4,9], for which the relevant exclusive branching ratios are of $O(10^{-5})$. Clean tests of the SM aim to see if the asymmetries for these decay modes are consistent with the allowed range for $\sin 2\beta$ and, moreover, whether they are equal to each other.

Among the channels given above, the most sensitive to the effect of the R-parity violating models are the ones that involve the second and the third family. Therefore we concentrate on these two modes: $b \rightarrow c\bar{c}s$ ($B_d \rightarrow \psi K_S$) and $b \rightarrow s\bar{s}s$ ($B_d \rightarrow \phi K_S$).

In the presence of new physics a non-negligible A_2 (with a non-trivial weak phase ϕ_2) could arise for both channels. In addition, there could be a new contribution to the mixing amplitude. Consequently, the asymmetries may measure angles that are different from β and, moreover, different from each other.

For simplicity we consider $\delta_1 = \delta_2$ for both channels, considering the fact that if they are not equal we have direct CP violation and our analysis can be generalized. It follows that the ratio between the conjugate amplitudes is a pure phase, $\bar{A}/A = e^{-2i\phi_D}$. From eq.(2.4) and eq.(2.5), we have:

$$\lambda = e^{-2i(\phi_M + \phi_D)}, \quad (3.2)$$

which implies that $|\lambda| = 1$ and consequently $a_{f_{CP}}^{\cos} = 0$.

In this case we can parameterize the new physics effects in $a_{f_{CP}}$ by writing

$$\lambda = e^{2i(\phi_0 + \delta\phi)}, \quad a_{f_{CP}}(t) = -\sin[2(\phi_0 + \delta\phi)] \sin(\Delta Mt), \quad (3.3)$$

where ϕ_0 is the phase predicted at leading order in the SM,

$$a_{f_{CP}}^{SM}(t) \approx -\sin[2\phi_0] \sin(\Delta Mt) \quad (3.4)$$

and $\delta\phi$ is the correction due to new physics (and subleading SM contributions, if they are not negligible).

The R-parity violating couplings can contribute to both the $B_d - \bar{B}_d$ mixing amplitude and the decay amplitude of a particular process. The effect in the mixing will translate into $\phi_M^{SM} \rightarrow \phi_M^{SM} + \delta\phi_M^R$ and this shift is the same for all B_d decays. Although it changes the value of the asymmetry with respect to the SM, it does not change the pattern predicted by the SM, which is that the CP asymmetries in $B_d \rightarrow \phi K_S$ and $B_d \rightarrow \psi K_S$ are equal. The effect in the decay, $\phi_D^{SM} \rightarrow \phi_D^{SM} + \delta\phi_D^R$, depends on the specific process. For the two channels we mentioned before we consider the difference between the angles $\phi(B_d \rightarrow f)$, obtained from the asymmetry measurement in the $B_d \rightarrow f$ decay, that has to be zero in the SM up to theoretical uncertainties [9],

$$|\phi(B_d \rightarrow \phi K_S) - \phi(B_d \rightarrow \psi K_S)| < 0.1. \quad (3.5)$$

This equation is likely to hold if there is not a large enhancement of the matrix element of the $b \rightarrow u\bar{u}s$ operator between the B_d initial state and the ϕK_S final state. In [9] the authors use $SU(3)$ relations to suggest some experimental tests that can constrain the size of this effect. These tests are likely to be done by the time that the CP measurements will start.

If eq.(3.5) holds, it makes sense to estimate the difference in the framework of SUSY models without R-parity and to check if it can be larger than the SM uncertainties.

IV. SUSY WITHOUT R-PARITY

In this section we present a short introduction to SUSY without R-parity. In contrast to the SM, in the Supersymmetric version of the SM, the general Lorentz and gauge invariant Lagrangian does not have the accidental symmetries of baryon number (B) and lepton number (L). The most general L and B violating superpotential is given by

$$W_R = \lambda_{ijk} L_i L_j \bar{l}_k + \lambda'_{ijk} L_i Q_j \bar{d}_k + \lambda'' \bar{u}_i \bar{d}_j \bar{d}_k + \mu_i L_i \phi_u; \quad (4.1)$$

where $i, j, k = 1, 2, 3$ are generation indices. $L_i (Q_i)$ are the lepton (quark) $SU(2)_L$ doublet superfields, $\bar{l}_j (\bar{d}_j, \bar{u}_j)$ are the charged leptons (down and up quark) $SU(2)_L$ singlet superfields and ϕ_u is the up Higgs superfield. The bilinear terms are relevant for the ν -masses [10,11], but they are not important for the present analysis so we will neglect them in the following. The first two terms are L -violating and the third is B -violating.

The combination of L and B violating operators will lead to rapid proton decay. Then an extra symmetry is required to enforce nucleon stability. Often an ad-hoc symmetry, called R-parity, is imposed to keep L and B symmetries intact. This symmetry assigns a charge $R = (-1)^{3B+L+2S}$ to each particle, where S is the particle's spin. Other possible choices are to impose only B or L .

For our analysis we consider SUSY extensions of the SM with highly suppressed B -violation ($\lambda'' = 0$), but without R-parity [12] and without lepton number [13] ($\lambda', \lambda \neq 0$), which represent interesting alternatives to the MSSM [14]. The λ' terms give rise to new contributions to B decays.

Non-leptonic B -decays are caused by b-quark transitions of the type $b \rightarrow q_1 \bar{q}_2 q_3$, with $q_1 \in \{d, s\}$ and $q_2, q_3 \in \{u, d, c, s\}$. The R-parity violating effective terms relevant for these kind of decays arise from the slepton mediated tree level diagrams. They are of the form:

$(\lambda'\lambda'/M_{(\tilde{l}\text{ or } \tilde{\nu})}^2)(\bar{q}_{(L,R)}b_{(R,L)})(\bar{q}_{(R,L)}q_{(L,R)})$, where $M_{\tilde{l}}$ and $M_{\tilde{\nu}}$ are the masses of the intermediate charged slepton and sneutrino. In the following analysis we neglect contributions of additional new physics operators which will affect the asymmetries at the loop level.

Several models that can explain the observed fermion mass hierarchy, like supersymmetric models with horizontal symmetries [15,16], also predict that R-parity violating couplings involving only the third and the second generations fields are the largest ones [13]. This is why we focus on $b \rightarrow c\bar{c}s$ and $b \rightarrow s\bar{s}s$ decays.

V. R-PARITY VIOLATING CONTRIBUTIONS TO THE DECAY AMPLITUDES

Referring to eq.(3.1), we can write the decay amplitude in the presence of R-parity violating couplings as:

$$A = A_{SM} + A_R = A_{SM}\Delta_D^R, \quad (5.1)$$

with

$$\Delta_D^R = 1 + r_D e^{i\theta_D}, \quad (5.2)$$

where $r_D = |A_R/A_{SM}|$ and $\theta_D = \arg(A_R/A_{SM})$. Note that r_D is often constrained phenomenologically while θ_D is not. Consequently the phase ϕ_D in the amplitude becomes $\phi_D = \phi_D^{SM} + \delta\phi_D$ where

$$\delta\phi_D = \arg(\Delta_D^R) = \arctan\left(\frac{r_D \sin \theta_D}{1 + r_D \cos \theta_D}\right). \quad (5.3)$$

From this equation we see that if $r_D \ll 1$ then $\delta\phi_D \leq r_D$. If however $r_D \geq 1$, then $\delta\phi_D$ could take any value, because it depends strongly on θ_D .

Since r_D plays an important role in our discussion, we next estimate its order of magnitude for the two processes.

A. Estimate of $r_D(B_d \rightarrow \phi K_S)$

The amplitude for this process is given by:

$$A(B_d \rightarrow \phi K_S) = \langle \phi K_S | \mathcal{H}_{eff} | B_d \rangle, \quad (5.4)$$

where

$$\mathcal{H}_{eff} = \mathcal{H}_{eff}^{SM} + \mathcal{H}_{eff}^R. \quad (5.5)$$

The Standard Model effective Hamiltonian relevant for this channel is:

$$\mathcal{H}_{eff}^{SM} = -\frac{G_F}{\sqrt{2}}(V_{ts}^* V_{tb}) \sum_{k=3}^{10} Q_k c_k(\mu), \quad (5.6)$$

where $c_k(\mu)$ are the Wilson coefficients, which are functions of the scale μ , with $\mu = O(m_b)$ being the relevant scale for b decays. The operators $Q_3..Q_6$ are QCD penguin operators and $Q_7..Q_{10}$ are electroweak penguin operators [17] .

The final ϕ requires that Q_k lead to a color-singlet term of the type $\bar{s}\gamma^\mu s$. When Fierz-transforming to obtain the required structure, some terms will be multiplied by $1/N_c$ (N_c is the number of colors).

Our calculation uses the method of Deshpande and He [18]. In particular, we use factorization to obtain the amplitude for exclusive processes. Using factorization introduces an unknown error of order one in the final result. We do not worry about uncertainties of this size because we are trying only to get order of magnitude estimates. In the SM we have:

$$A^{SM}(B_d \rightarrow \phi K_S) = -\frac{G_F}{\sqrt{2}}(V_{ts}^* V_{tb})C \langle K_S | \bar{s}\gamma^\mu (1 - \gamma_5) b | B_d \rangle \langle \phi | \bar{s}\gamma^\mu s | 0 \rangle. \quad (5.7)$$

C is the following combination of Wilson coefficients:

$$C = (1 + \frac{1}{N_c}) c_3 + (1 + \frac{1}{N_c}) c_4 + c_5 + \frac{1}{N_c} c_6 \\ - \frac{1}{2}(c_7 + \frac{1}{N_c} c_8 + (1 + \frac{1}{N_c}) c_9 + (1 + \frac{1}{N_c}) c_{10}). \quad (5.8)$$

Using the results of ref. [17], we can give an estimate of C in the leading logarithmic approximation (LO), $C \simeq 2 \times 10^{-2}$.

For $b \rightarrow s\bar{s}s$ transitions the R-parity violating contributions come from the sneutrino mediated diagrams which give rise to these new terms:

$$c_1^R Q_1^R = \frac{1}{M_{\tilde{\nu}}^2} (\lambda_{i23}^* \lambda'_{i22}) (\bar{s}_L^\alpha b_R^\alpha) (\bar{s}_R^\beta s_L^\beta) \\ c_2^R Q_2^R = \frac{1}{M_{\tilde{\nu}}^2} (\lambda'_{i32} \lambda_{i22}^*) (\bar{s}_R^\alpha b_L^\alpha) (\bar{s}_L^\beta s_R^\beta). \quad (5.9)$$

The Fierz transformation introduces color suppression factors:

$$Q_1^R \Rightarrow \frac{1}{2} \frac{1}{N_c} (\bar{s}_L \gamma^\mu s_L) (\bar{s}_R \gamma^\mu b_R) \\ Q_2^R \Rightarrow \frac{1}{2} \frac{1}{N_c} (\bar{s}_R \gamma^\mu s_R) (\bar{s}_L \gamma^\mu b_L). \quad (5.10)$$

For $\mathcal{H}_{eff}^R$ we find:

$$\mathcal{H}_{eff}^R = \frac{1}{M_{\tilde{\nu}}^2} \frac{1}{N_c} \frac{1}{8} [(\lambda_{i23}^* \lambda'_{i22}) (\bar{s}\gamma^\mu (1 - \gamma_5) s) (\bar{s}\gamma^\mu (1 + \gamma_5) b) \\ + (\lambda'_{i32} \lambda_{i22}^*) (\bar{s}\gamma^\mu (1 + \gamma_5) s) (\bar{s}\gamma^\mu (1 - \gamma_5) b)]. \quad (5.11)$$

Since

$$\langle K_S | \bar{s}\gamma^\mu (1 - \gamma_5) b | B_d \rangle = \langle K_S | \bar{s}\gamma^\mu (1 + \gamma_5) b | B_d \rangle = \langle K_S | \bar{s}\gamma^\mu b | B_d \rangle, \quad (5.12)$$

and

$$\langle \phi | \bar{s} \gamma^\mu (1 - \gamma_5) s | 0 \rangle = \langle \phi | \bar{s} \gamma^\mu (1 + \gamma_5) s | 0 \rangle = \langle \phi | \bar{s} \gamma^\mu s | 0 \rangle, \quad (5.13)$$

the R-parity contribution to the amplitude modifies the SM amplitude by an overall coefficient:

$$\begin{aligned} A(B_d \rightarrow \phi K_S) &= A^{SM}(B_d \rightarrow \phi K_S) + A^R(B_d \rightarrow \phi K_S) \\ &= \left[-\frac{G_F}{\sqrt{2}} (V_{ts}^* V_{tb}) C + \frac{1}{M_{\tilde{\nu}}^2} \frac{1}{N_c} \frac{1}{8} (\lambda_{i23}'^* \lambda_{i22}' + \lambda_{i32}' \lambda_{i22}'^*) \right] \langle \phi | \bar{s} \gamma^\mu s | 0 \rangle \langle K_S | \bar{s} \gamma^\mu b | B_d \rangle. \end{aligned} \quad (5.14)$$

We are now in a position to give $r_D(B_d \rightarrow \phi K_S)$:

$$r_D(B_d \rightarrow \phi K_S) = \frac{|\frac{1}{M_{\tilde{\nu}}^2} \frac{1}{N_c} \frac{1}{8} (\lambda_{i23}'^* \lambda_{i22}' + \lambda_{i32}' \lambda_{i22}'^*)|}{|\frac{g^2}{8M_W^2} (V_{ts}^* V_{tb}) C|}. \quad (5.15)$$

Putting $|V_{ts}^* V_{tb}| = 4 \times 10^{-2}$, $N_c = 3$, $C \simeq 2 \times 10^{-2}$ we estimate

$$r_D \simeq 8 \times 10^2 |\lambda_{i23}'^* \lambda_{i22}' + \lambda_{i32}' \lambda_{i22}'^*| \left(\frac{M_W}{M_{\tilde{\nu}}} \right)^2. \quad (5.16)$$

The bound $BR(b \rightarrow X_s \nu \bar{\nu}) \leq 7.7 \times 10^{-4}$ [19] implies the following limits on products of R-parity violating couplings [20]:

$$\lambda_{i32}' \lambda_{i22}'^* < 1.5 \times 10^{-3} \left(\frac{m_{\tilde{d}_L}}{100 \text{ GeV}} \right)^2. \quad (5.17)$$

For the product of the other couplings we use the limits found in [21]

$$\lambda_{i23}'^* \lambda_{i22}' < 1.4 \times 10^{-4} \left(\frac{m_{\tilde{d}_R}}{100 \text{ GeV}} \right)^2. \quad (5.18)$$

Using these limits and taking $m_{\tilde{d}} \sim M_{\tilde{\nu}}$ we get

$$r_D(B_d \rightarrow \phi K_S) \leq 0.8, \quad (5.19)$$

and so

$$\delta\phi(B_d \rightarrow \phi K_S) \lesssim 0.8. \quad (5.20)$$

B. Estimate of $r_D(B_d \rightarrow \psi K_S)$

In [5] the estimate of the maximal value of r_D was done considering the matrix elements of the R-parity violating and SM operators of the same order. We obtain the same result following the steps of the previous analysis for

$$A(B_d \rightarrow \psi K_S) = \langle \psi K_S | \mathcal{H}_{eff} | B_d \rangle. \quad (5.21)$$

The relevant SM Hamiltonian is:

$$\mathcal{H}_{eff}^{SM} = \frac{G_F}{\sqrt{2}} (V_{cb}^* V_{cs}) \sum_{k=1}^{10} Q_k c_k(\mu). \quad (5.22)$$

We find:

$$A^{SM}(B_d \rightarrow \psi K_S) = \frac{G_F}{\sqrt{2}} (V_{cb}^* V_{cs}) C' \langle K_S | \bar{s} \gamma^\mu (1 - \gamma_5) b | B_d \rangle \langle \psi | \bar{c} \gamma^\mu c | 0 \rangle, \quad (5.23)$$

where $C' = c_1 + \frac{1}{N_c} c_2$. In the LO, $C' \simeq 8 \times 10^{-2}$ [17].

For $b \rightarrow c \bar{c} s$ transitions the R-parity violating contributions come from the charged slepton mediated diagrams which give rise to a new term which Fierz transforms to:

$$c_1^R Q_1^R = \frac{1}{M_l^2} \frac{1}{N_c} \frac{1}{8} (\lambda_{i23}'^* \lambda_{i22}') (\bar{c}_L \gamma^\mu c_L) (\bar{s}_R \gamma^\mu b_R). \quad (5.24)$$

The R-parity contribution to the amplitude is to modify the overall coefficient:

$$\begin{aligned} A(B_d \rightarrow \psi K_S) &= A^{SM}(B_d \rightarrow \psi K_S) + A^R(B_d \rightarrow \psi K_S) \\ &= \left[\frac{G_F}{\sqrt{2}} (V_{cb}^* V_{cs}) C' + \frac{1}{M_l^2} \frac{1}{N_c} \frac{1}{8} (\lambda_{i23}'^* \lambda_{i22}') \right] \\ &\quad \langle \psi | \bar{c} \gamma^\mu c | 0 \rangle \langle K_S | \bar{s} \gamma^\mu b | B \rangle, \end{aligned} \quad (5.25)$$

leading to

$$r_D(B_d \rightarrow \psi K_S) = \frac{|\frac{1}{M_l^2} \frac{1}{N_c} \frac{1}{8} (\lambda_{i23}'^* \lambda_{i22}')|}{|\frac{g^2}{8M_W^2} (V_{cb}^* V_{cs}) C'|} \simeq 2 \times 10^2 |\lambda_{i23}'^* \lambda_{i22}'| \left(\frac{M_W}{M_l} \right)^2. \quad (5.26)$$

Putting the same limits as before for $\lambda_{i23}'^* \lambda_{i22}'$, and taking $M_l \simeq m_{\tilde{d}}$, we find $r_D \leq 0.02$ which is in agreement with what found in [5]. Then we have

$$\delta\phi(B_d \rightarrow \psi K_S) \lesssim 0.02. \quad (5.27)$$

We conclude that in the presence of R-parity violating couplings it is possible to obtain:

$$|\phi(B_d \rightarrow \phi K_S) - \phi(B_d \rightarrow \psi K_S)| \sim O(1). \quad (5.28)$$

Comparing this result with eq.(3.5) we learn that new physics in decay amplitudes could lead to deviation from the pattern of CP violation in B_d decays larger than the SM uncertainties.

VI. EFFECTS OF R-PARITY VIOLATING COUPLINGS ON $B_D - \bar{B}_D$ MIXING

In order to give a complete analysis of how R parity violating couplings affect the CP asymmetries, we have to consider the effect on $B_d - \bar{B}_d$ mixing.

The mixing amplitude can be written as

$$M_{12} = M_{12}^{SM} + M_{12}^R = M_{12}^{SM} \Delta_M^R, \quad (6.1)$$

where we define

$$\Delta_M^R = 1 + r_M e^{i\theta_M}, \quad (6.2)$$

with $r_M = |M_{12}^R/M_{12}^{SM}|$ and $\theta_M = \arg[\lambda_{i13}^* \lambda'_{i31}/(V_{tb}^* V_{td})^2]$. The mixing angle becomes $\phi_M = \phi_M^{SM} + \delta\phi_M$, where

$$\delta\phi_M = \arg(\Delta_M^R) = \arctan\left(\frac{r_M \sin \theta_M}{1 + r_M \cos \theta_M}\right). \quad (6.3)$$

In order to estimate the contribution to the mixing, we first study the phenomenological constraints on r_M and then we estimate it in the model with horizontal symmetry.

Note that the only experimental limits we have on the R-parity violating couplings that contribute to the mixing come from the bound on ΔM_{B_d} itself. Since the R-parity violating contribution could saturate ΔM_B and the SM contribution could be less than a half of it, it is clear that $r_M > 1$ is allowed.

From eq.(6.3) we see that, if $r_M > 1$, $\delta\phi_M$ depends strongly on θ_M , which, in turn, depends on the new phases of the λ' couplings. As there are no experimental constraints on the value of these phases, the asymmetries could have any value, unlike the SM case where the phases are constrained, i.e. $0.3 \lesssim \sin 2\beta \lesssim 0.9$.

Although we concluded that r_M could be large, it is still useful to give an explicit expression for it. This is done in the next subsection.

Estimate of $r_M(B_d \rightarrow f)$

The mixing matrix element is given by:

$$|M_{12}| = \frac{1}{M_{B_d}} |\langle \bar{B}_d | \mathcal{H}_{eff}(\Delta B = 2) | B_d \rangle|, \quad (6.4)$$

the SM contribution is [17]

$$M_{12}^{SM}(B_d) = \frac{G_F^2}{12\pi^2} \eta_B M_{B_d} (B_{B_d} f_{B_d}^2) M_W^2 S_0(x_t) (V_{tb}^* V_{td})^2, \quad (6.5)$$

where $\eta_B = 0.55$, [22], $x_t = (m_t/M_W)^2$ and $S_0(x_t)$ is given in [23].

R-parity violating terms contribute to M_{12} at tree level through sneutrino-exchange. In fact, the effective Hamiltonian term for $B_d - \bar{B}_d$ is

$$\mathcal{H}_{eff}^R = \frac{\lambda_{i13}^* \lambda'_{i31}}{M_{\tilde{\nu}}^2} Q_S(\Delta B = 2) + h.c. \quad (6.6)$$

where

$$Q_S(\Delta B = 2) = (\bar{d}_L b_R)(\bar{d}_R b_L) \quad (6.7)$$

In the vacuum insertion approximation [24]:

$$\langle \bar{B}_d | Q_S(\Delta B = 2) | B_d \rangle = \left(\frac{M_{B_d}^2}{(m_b + m_d)^2} + \frac{1}{6} \right) f_{B_d}^2 M_{B_d}^2. \quad (6.8)$$

Then the R-parity violating mixing term is

$$M_{12}^R = \frac{\lambda'_{i13} \lambda'_{i31}}{M_{\tilde{\nu}}^2} \left(\frac{M_{B_d}^2}{(m_b + m_d)^2} + \frac{1}{6} \right) f_{B_d}^2 M_{B_d}, \quad (6.9)$$

leading to

$$r_M = \frac{\left| \frac{\lambda'_{i13} \lambda'_{i31}}{M_{\tilde{\nu}}^2} \left(\frac{M_{B_d}^2}{(m_b + m_d)^2} + \frac{1}{6} \right) \right|}{\left| \frac{G_F^2}{12\pi^2} \eta_B B_{B_d} M_W^2 S_0(x_t) (V_{tb}^* V_{td})^2 \right|} \simeq 10^8 |\lambda'_{i13} \lambda'_{i31}| \left(\frac{100 \text{ GeV}}{M_{\tilde{\nu}}} \right)^2. \quad (6.10)$$

VII. R-PARITY VIOLATION IN THE FRAMEWORK OF HORIZONTAL SYMMETRIES

In the previous sections we found the upper bounds on $r_D(B_d \rightarrow f)$ from phenomenological constraints and we gave the general expression of r_M as a function of the R-parity violating couplings. In this section we estimate r_D and r_M in the framework of supersymmetric models with Abelian horizontal symmetries that have been thoroughly investigated in [15,16]. These models explain naturally the order of magnitude of the fermion masses, the hierarchy among them and the CKM angles. Assuming a horizontal $U(1)$ symmetry with a small breaking parameter, $\varepsilon \sim 0.2$, allows us to estimate the size of the $\lambda' L$ violating couplings relevant for our processes and to work out numerical predictions for r_D . Most of the L -violating couplings are suppressed with respect to the corresponding Yukawa couplings Y_{ij} . They can be estimated as [7]

$$\lambda'_{kij} \sim Y_{ij}^d \varepsilon^{H(L_k) - H(\Phi_d)}, \quad (7.1)$$

where $H(\psi)$ is the charge assigned to the field ψ . Using the equations (2.2)-(2.4) of ref. [15] we obtain

$$\lambda'_{kij} \lesssim \sqrt{G_F \tan \beta} (|V_{ij}|)^{\text{sign}(j-i)} m_{d_j} \varepsilon^{H(L_k) - H(\Phi_d)}. \quad (7.2)$$

The equation (7.2) shows that like the lepton and down quark Yukawa couplings, the λ' couplings increase with $\tan \beta = \langle \phi_u \rangle / \langle \phi_d \rangle$.

A. Estimate of r_D for the two processes

We consider two different frameworks of the horizontal symmetries that allow us to find order of magnitude estimates for r_D in these models.

a. In a framework where the neutrino masses are suppressed only by the horizontal symmetries [13] we can write

$$\varepsilon^{H(L_k) - H(\Phi_d)} \lesssim \sqrt{\frac{m_{\nu_\tau}}{m_Z}}, \quad (7.3)$$

therefore the constraints on the λ' couplings come from the bound on the mass of the τ -neutrino, $m_{\nu_\tau} \lesssim 18 \text{ MeV}$ [25], obtained from direct experiments.

The suppression of the λ' couplings is then related to fermion parameters in the following way:

$$\begin{aligned} r_D(B_d \rightarrow \phi K_S) &\lesssim 2 \times 10^{-3} G_F \tan^2 \beta \frac{m_s^2}{|V_{cb} V_{ts}|} \\ r_D(B_d \rightarrow \psi K_S) &\lesssim 5 \times 10^{-4} G_F \tan^2 \beta m_s m_b. \end{aligned} \quad (7.4)$$

We fix $\tan \beta = 1$ and obtain the numerical results:

$$\begin{aligned} r_D(B_d \rightarrow \phi K_S) &\lesssim 5 \times 10^{-7} \\ r_D(B_d \rightarrow \psi K_S) &\lesssim 5 \times 10^{-9}, \end{aligned} \quad (7.5)$$

leading to

$$|\phi(B_d \rightarrow \phi K_S) - \phi(B_d \rightarrow \psi K_S)| \lesssim 5 \times 10^{-7}. \quad (7.6)$$

Then in this framework we obtain unobservable deviations from the SM predictions.

b. Within a framework in which the neutrino masses are not suppressed by the horizontal symmetries but by alternative mechanisms, (like the alignment of the μ and B terms [10]) we relate our couplings to the ones involved in processes where the R-parity violating terms induce flavour changing neutral currents, like $B - \bar{B}$ and $K - \bar{K}$ mixing, which are the most constrained. In this framework we get the maximal values allowed for λ' in agreement with all present constraints.

The strongest bound is the one on the product of the couplings involved in $K - \bar{K}$ mixing [26]

$$\lambda'_{i21} \lambda'_{i12} \leq 5 \times 10^{-9}. \quad (7.7)$$

Then writing our couplings in terms of these ones and of the fermion parameters we obtain for r_D of the two processes:

$$\begin{aligned} r_D(B_d \rightarrow \phi K_S) &\sim 20 \left(\frac{|V_{us}|}{|V_{cd} V_{cb} V_{ts}|} \frac{m_s}{m_d} \right) |\lambda'_{i21} \lambda'_{i12}| \\ r_D(B_d \rightarrow \psi K_S) &\sim 5 \left(\frac{|V_{us}|}{|V_{cd}|} \frac{m_b}{m_d} \right) |\lambda'_{i21} \lambda'_{i12}|. \end{aligned} \quad (7.8)$$

Putting the numerical values and the constraints on the couplings, we have:

$$\begin{aligned} r_D(B_d \rightarrow \phi K_S) &\lesssim 1 \times 10^{-3} \\ r_D(B_d \rightarrow \psi K_S) &\lesssim 1 \times 10^{-5}, \end{aligned} \quad (7.9)$$

leading to

$$|\phi(B_d \rightarrow \phi K_S) - \phi(B_d \rightarrow \psi K_S)| \lesssim 1 \times 10^{-3}. \quad (7.10)$$

We see that also in this case we do not obtain sizeable effects.

From eq.(7.6) and eq.(7.10) we conclude that this type of R-parity violating SUSY models are unlikely to be signaled out through the comparison of these two CP asymmetries. However in [7] it was found that a signature of these models could indeed be detected in rare B decays in the third family leptons. Therefore we can conclude that rare B decays are more sensitive to this kind of effects than CP asymmetries.

B. Estimate of r_M

For the mixing, in the framework *a* of the horizontal symmetries we have:

$$r_M \lesssim 10^4 G_F \tan^2 \beta \frac{|V_{ub}|}{|V_{tb}|} m_b m_d. \quad (7.11)$$

Inserting the numerical values we obtain:

$$r_M \lesssim 4 \times 10^{-3}, \quad (7.12)$$

and the correction to the mixing phase is limited to be:

$$\delta\phi_M \lesssim 4 \times 10^{-3}. \quad (7.13)$$

We conclude also that this effect is too small to be signaled out through the measurement of the CP asymmetries.

In the framework *b* we find for r_M

$$r_M \sim 10^8 \left(\frac{|V_{ub}|}{|V_{ts}V_{cd}|} \frac{m_b}{m_s} \right) |\lambda'_{i21} \lambda'_{i12}|. \quad (7.14)$$

Putting the numerical values and the constraint on the product of the couplings, we obtain

$$r_M \sim 1. \quad (7.15)$$

Then $\delta\phi_M$ depends strongly on θ_M ,

$$\delta\phi_M \sim \frac{\theta_M}{2}, \quad (7.16)$$

on which we do not have any constrain. This implies that unlike the SM case the CP asymmetries can assume any value. Even if in this case the new physics can affect strongly the mixing phase, again a measurement of the CP asymmetries does not allow us to identify this particular kind of new physics.

VIII. CONCLUSIONS

We have studied new CP violating effects that can arise in models without R-parity and without lepton number. We focused on the CP asymmetries for the two channels $B_d \rightarrow \psi K_S$ and $B_d \rightarrow \phi K_S$ because these observables are experimentally accessible, have small theoretical uncertainties and are sensitive to this type of new physics.

In the SM these decays measure directly the phase β , up to small corrections due to the presence of subleading contributions of order 10% in $B_d \rightarrow \phi K_S$.

In general we have found that $B_d \rightarrow \phi K_S$ and the $B-\bar{B}$ mixing may be strongly affected, while the amplitude of the mode $B_d \rightarrow \psi K_S$ remains almost the same that in the SM. The result of the R-parity violating couplings could be a deviation of both the CP asymmetries from $\sin 2\beta$ and in a sizeable difference between them. In SUSY models without R-parity

and with Abelian horizontal symmetries we have considered two possible situations. When ν -masses are suppressed only by the horizontal symmetry, we have found negligible effects in the amplitudes and a weak effect in the mixing. In models where the μ and the B terms are aligned so that ν masses arise only from loop effects (or nonrenormalizable terms), the effect of the R-parity violating couplings could be larger. However, also in this case the effect on the amplitudes remains small while the effect on the mixing can be sizeable. We conclude that the CP asymmetries are not sensitive to this kind of new physics. Other observables like branching ratios of rare B decays into third family leptons are better suited for the study of this kind of effects [7].

ACKNOWLEDGMENTS

I thank Yossi Nir for suggesting this work, Roberto Casalbuoni, Yuval Grossman for relevant contributions to the paper and Adam Falk, Alex Kagan, Enrico Nardi for useful comments on the manuscript. I also acknowledge the Particle Theory Group of the Weizmann Institute of Science, where this research was carried out, for the hospitality and for the pleasant working atmosphere.

REFERENCES

- [1] T. Nakada, preprint PSI-PR-96-22, hep-ex/9609015;
F. Grancagnolo, Italy INFN-INFN-AE-90-07, Les Arcs 1990, Proceedings, High energy hadron interactions, 249-254;
CLEO Collaboration, CLNS-94-1277, Feb 1994. 146pp.;
J.N. Butler, FERMILAB-PUB-95-363, Nucl. Instrum. Meth A368, 145-151, (1995).
- [2] For a review see e.g. Y. Nir, Lectures presented in the 20th SLAC Summer Institute, SLAC-PUB-5874 (1992);
I. Dunietz, Ann. Phys. **184** (1988) 350;
Review of Particle Properties, Phys. Rev. **D 54** (1996) 507.
- [3] Y. Nir and H.R. Quinn, Ann. Rev. Nucl. Part. Sci. **42** (1992) 211;
M. Gronau and D. London, Phys. Rev. **D 55** (1997) 2845;
Y. Nir, Y. Grossman and R. Rattazzi, SLAC-PUB-7379, hep-ph/9701231;
R. Fleisher, CERN-TH/97-241, hep-ph/9709291;
Y. Nir, WIS-97/28/Sep-PH, hep-ph/9709301;
M. Ciuchini *et al.*, Phys. Rev. Lett. **79** (1997) 978;
D. London and A. Soni, Phys. Lett. **B 407** (1997) 61;
R. Barbieri and A. Strumia, Nucl. Phys. **B 508** (1997) 3.
- [4] Y. Grossman and Mihir P. Worah, Phys. Lett. **B 395** (1997) 241.
- [5] D.E. Kaplan, hep-ph/9703347.
- [6] S. A. Abel, Phys. Lett. **B 410** (1997) 173.
- [7] D. Guetta and E. Nardi, Phys. Rev. **D 58** 012001 (1998).
- [8] See e.g. T. Kurimoto, hep-ph/9706221.
- [9] Y. Grossman, G. Isidori and M. P. Worah, SLAC-PUB-7614 hep-ph/9708305.
- [10] E. Nardi, Phys. Rev. **D 55** (1997) 5772.
- [11] R. Hempfling, Nucl. Phys. **B 478** (1996) 3;
B. de Carlos and P.L. White, Phys. Rev. **D 54** (1996) 3427;
H.P. Nilles and N. Polonsky, Nucl. Phys. **B 484** (1997) 33.
- [12] C.S. Aulakh and R.N. Mohapatra, Phys. Lett. **B 119** (1982) 136;
I.-H. Lee Phys. Lett. **B 138** (1984) 121; Nucl. Phys. **B 246** (1984) 120;
S. Dawson, Nucl. Phys. **B 261** (1985) 297;
G. Ross and J. Valle, Phys. Lett. **B 151** (1985) 375;
J. Ellis *et al.*, Phys. Lett. **B 150** (1985) 142;
R. N. Mohapatra, Phys. Rev. **D 34** (1986) 3457;
A. Santamaria and J.W.F. Valle, Phys. Lett. **B 195** (1987) 423;
S. Dimopoulos and L.J. Hall, Phys. Lett. **B 207** (1987) 210;
V. Barger, G.F. Giudice, and T. Han, Phys. Rev. **D 40** (1989) 2987;
D.E. Brahm, L.J. Hall, and S. Hsu, Phys. Rev. **D 42** (1990) 1860;
K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **64** (1990) 1705;
R. Barbieri, M. Guzzo, A. Masiero, and D. Tommasini, Phys. Lett. **B 252** (1990) 251;
E. Roulet and D. Tommasini, Phys. Lett. **B 256** (1991) 218;
K. Enkvist, A. Masiero, and A. Riotto, Nucl. Phys. **B 373** (1992) 95;
J.C. Romao and J.W.F. Valle, Nucl. Phys. **B 381** (1992) 87;
R.M. Godbole, P. Roy, and X. Tata, Nucl. Phys. **B 401** (1993) 67;
I. Hinchliffe and T. Kaeding, Phys. Rev. **D 47** (1993) 279;

- K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **75** (1995) 2276;
A.S. Joshipura and M. Nowakowski, Phys. Rev. **D 51** (1995) 2421;
F. de Campos *et. al.*, Nucl. Phys. **B 451** (1995) 3;
M.Hirsch, H. Klapdor-Kleingrothaus and S. Kovalenko, Phys. Rev. Lett. **75** (1995) 17;
G. Bhattacharyya, J. Ellis and K. Sridhar, Mod. Phys. Rev. Lett. A **10** (1995) 1583;
S. Roy and B. Mukhopadhyaya, Phys. Rev. **D 55** (1997) 7020.
- [13] T. Banks, Y. Grossman, E. Nardi and Y. Nir, Phys. Rev. **D 52** (1995) 5319.
 - [14] For reviews on supersymmetry and the MSSM, see *Review of Particles Properties*, Phys. Rev. **D 54** (1996) 687, and references within.
 - [15] M. Leurer, Y. Nir and N. Seiberg, Nucl. Phys. **B 398** (1993) 319.
 - [16] Y. Nir and N. Seiberg, Phys. Lett. **B 309** (1993) 337;
M. Leurer, Y. Nir and N. Seiberg, Nucl. Phys. **B 420** (1994) 468.
 - [17] A.J. Buras and R. Fleischer hep-ph/9704376.
 - [18] N.G.Deshpande and Xiao-Gang He, Phys. Lett. **B 336** (1994) 471.
 - [19] ALEPH Collaboration, Report no. PA10-019, presented at the “28th International Conference on High Energy Physics”, Warsaw, Poland (unpublished).
 - [20] Y. Grossman, Z. Ligeti, E. Nardi Nucl. Phys. **B 465** (1996) 369-398; Erratum Nucl. Phys. **B 480** (1996) 753.
 - [21] K. Agashe and M.Graesser, Phys. Rev. **D 54** (1996) 4445.
 - [22] A.J.Buras, M. Jamin and P.H. Weisz, Nucl. Phys. **B 347** (1990) 3421.
 - [23] T.Inami and C.S. Lim, Prog. Theor. Phys. **65** (1981) 297.
 - [24] G.Ecker and W.Grimus, Nucl. Phys. **B 258** (1985) 328.
 - [25] G. Fiorillo *et al.*, Proceeding Nucl. Phys. **B**, Proc. Suppl. **66** (1998).
 - [26] D.Choudhury and P.Roy, Phys. Lett. **B 378** (1996) 153.